



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

MISCELLANEOUS.

155. Proposed by A. H. HOLMES, Brunswick, Maine.

There are two vessels, one containing a gallons of alcohol, the other b gallons of water. Suppose that c gallons are simultaneously taken from each and poured into the other, how many times must this be done so that there will be the same proportion of alcohol to water in each vessel?

156. Proposed by R. D. CARMICHAEL, Hartselle, Ala.

There exist no multiply perfect odd numbers of multiplicity n containing only n distinct primes.

UNSOLVED PROBLEMS.

NOTE. The following problems still remain unsolved (in our columns).

Average and Probability, 164 Proposed by J. O. MAHONEY, B.E., M.Sc., Central High School, Dallas, Tex.

If m is prime, and the numbers 0, 1, 2, ..., $m^2 - 1$ are placed at random in the form of a square, the probability that the square is hyper-magic is $(m-1)m/(m^2-2)!$

Algebra, 179. Proposed by DR. L. E. DICKSON, The University of Chicago.

Find the roots of the algebraically solvable quintic equation

$$x^5 + qx^2 + px + \frac{1}{8} \left[\frac{q^2}{p} - \frac{p^3}{5q} \right] = 0.$$

NOTES AND NEWS.

Mr. B. M. Rastall is assistant in mathematics at the University of Wisconsin.

Mr. E. A. Moritz, Mr. R. S. Peatter, and Mr. E. R. Smith have been appointed instructors in mathematics at the University of Wisconsin.

The University of North Carolina has granted a years leave of absence to Mr. M. H. Stacy, who will pursue a graduate course in mathematics at Cornell University.

The American Association for the Advancement of Science held its annual meeting during the holidays at New Orleans, La. Section A: Mathematics and Astronomy, was presided over by Dr. W. S. Eichelberger, of the United States Naval Observatory at Washington, D. C. Professor L. G. Weld, of the University of Iowa, is secretary of the Section.